

## Vaccination schedule for Pet animals

Sl.no	Name of disease	Vaccine	Age at first dose	Booster dose	Subsequent dose	Dose	Route
1	Canine Distemper, Canine Hepatitis, Parvoviral enteritis & Parainfluenza	Live attenuated vaccine	6-8 weeks	10-12 weeks	Annually	1 ml	S/C or I/M
2	Rabies (Prophylactic)	Inactivated virus adjuvanted in Gel	3 months and above	-	Annually	1 ml	S/C or I/M
	Rabies (Post-bite)	Inactivated virus adjuvanted in Gel	Zero day (Immediately after suspected bite)	3 <sup>rd</sup> day	7 <sup>th</sup> , 14 <sup>th</sup> , 28 <sup>th</sup> & 90 <sup>th</sup> day after first dose	1 ml	S/C or I/M
3	Feline panleucopenia, Feline rhinotracheitis & Feline calicivirus infections	Modified live vaccine	9 weeks	12-13 weeks	Annually	1 ml	S/C

### Points to be noted:

1. Animals should be in good health at the time of vaccination. Do not vaccinate animals which are already in stress.
2. Try to deworm animals & remove ecto-parasites one to two weeks prior to vaccination.
3. Strictly follow vaccination schedule after consultation with veterinary experts.
4. Keep records of vaccination.
5. Maintain cold chain for vaccines which may have to keep for 2-8 degree Celsius temperature.
6. Create stress free environment for animals after vaccination.
7. It is recommended to follow manufacturer's instructions.